## **\* Immuno**Reagents,Inc.

6003 Chapel Hill Road Suite 153 Raleigh, NC 27607

## **Product Specifications**

Product: Description: Part Number: Concentration: Amount: Conjugate:	Chicken anti-Rat IgG (H&L) - Affinity Pure Chicken anti-Rat IgG (H&L) - Affinity Pure, TRITC Conjugate CkxRt-003-DRHOX 1.0 mg/ml (E 1% at 280 nm = 13.2) 1.0 mg Tetramethylrhodamine-5-isothiocyanate (TRITC) Amax = 550 nm; Emax = 570 nm
Form:	Lyophilized
Purification:	Affinity purified using solid phase Rat IgG
Purity:	Affinity purified antibody is > 95% based on SDS-PAGE
Host:	Chicken
Immunogen:	Purified Rat IgG, whole molecule
Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v)
	BSA, Protease/IgG free
Preservative:	0.05% (w/v) Sodium Azide
Reconstitution:	Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.
Storage:	2-8 °C
Shelf Life:	1 year from date of receipt. Prepare working dilution prior to use and then discard.
Specificity:	Based on IEP, this antibody reacts with:
	<ul> <li>heavy (γ) chains on rat IgG</li> </ul>
	<ul> <li>light chains on all rat immunoglobulins</li> </ul>
Cross Reactivity:	Based on IEP, no reactivity is observed to:
	<ul> <li>non-immunoglobulin rat serum proteins</li> </ul>
Country of Origin:	Chicken Ig fraction was prepared using eggs from healthy hens of US origin.
Applications:	Flow Cytometry
	Immunofluorescence
	ELISA
	Immunomicroscopy

Fax: 919-831-2240

Disclaimer:

For *in vitro* Laboratory Use Only. Not for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license under any patent of ImmunoReagents, Inc. **Product may not be resold or modified for resale without prior written approval of ImmunoReagents, Inc.**