## **\* Immuno**Reagents,Inc.

6003 Chapel Hill Road Suite 153 Raleigh, NC 27607

## **Product Specifications**

| Product:<br>Description: | Donkey anti-Mouse IgG (H&L), F(ab)'2 Fragment<br>Donkey anti-Mouse IgG (H&L) - F(ab)'2 Fragment, DyLight®800<br>Conjugate, min x w/bovine, chicken, goat, guinea pig, hamster, horse,<br>human, rabbit, rat or sheep IgG |
|--------------------------|--|
| Part Number:             | DkxMu-003-L800NHSX   |
| Concentration:           | 1 mg/ml (E 1% at 280 nm = 13.0)  |
| Amount:                  | 0.5 mg   |
| Conjugate:               | DyLight® 800 (Ex = 770 nm; Em = 794 nm)  |
| Form:                    | Lyophilized  |
| Purification:            | Affinity purified using solid phase Mouse IgG  |
| Purity:                  | > 90% based on SDS-PAGE Small amounts of intact IgG may be present.  |
| Host:                    | Donkey   |
| Immunogen:               | Purified Mouse IgG, whole molecule   |
| Buffer:                  | 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v)  |
|                          | BSA, Protease/IgG free   |
| Preservative:            | 0.05% (w/v) Sodium Azide   |
| Reconstitution:          | Rehydrate with 0.55 ml of deionized water and let stand 30 minutes at  |
|                          | room temperature to dissolve. (Product has been overfilled to ensure   |
|                          | complete recovery.) Centrifuge to remove any particulates. Prepare fresh   |
|                          | working dilution daily.  |
| Storage:                 | Store freeze-dried powder at 2-8 °C.   |
| Shelf Life:              | Product is stable for up to 4 weeks at 2-8°C after rehydration. For  |
|                          | extended storage after rehydration, add an equal volume of glycerol and  |
|                          | store at -20°C.  |
| Specificity:             | Based on IEP, this antibody reacts with:   |
|                          | <ul> <li>heavy (γ) chains on mouse IgG</li> </ul>  |
|                          | <ul> <li>light chains on all mouse immunoglobulins</li> </ul>  |
|                          |  |
| Cross Reactivity:        | Based on IEP, no reactivity is observed to:  |
|                          | - non immunoglobulin mouse corum immunoglobuling   |
|                          | <ul> <li>non-immunoglobulin mouse serum immunoglobulins</li> <li>IgG from bovine, chicken, goat, guinea pig, hamster, horse, human,</li> </ul>   |
|                          | rabbit, rat or sheep   |
|                          |  |
| Country of Origin:       | Donkey serum was obtained from healthy animals of US origin and under  |
|                          | the care of a registered veterinarian.   |
|                          |  |
| Phone: 919-831-2240      | Fax: 919-831-2240 email: info@immunoreagents.com   |

| Applications: | Flow Cytometry  |
|---------------|---|
|               | Immunofluorescence  |
|               | ELISA   |
|               | Immunomicroscopy  |
| Disclaimer:   | For <i>in vitro</i> Laboratory Use Only. Not for diagnostic or therapeutic use. Not<br>for human or animal consumption. Suggested applications of our products<br>are not recommendations to use our products in violation of any patent or<br>as a license under any patent of ImmunoReagents, Inc. <b>Product may not</b><br><b>be resold or modified for resale without prior written approval of</b><br><b>ImmunoReagents, Inc.</b> |
|               |   |
| Trademark:    | DyLight® is a trademark of Thermo Fisher Scientific, Inc. and its subsidiaries.   |