*** Immuno**Reagents,Inc.

6003 Chapel Hill Road Suite 153 Raleigh, NC 27607

Product Specifications

Product: Description:	Goat anti-Human IgA (α chain), F(ab)'2 Fragment - Affinity Pure Goat anti-Human IgA (α chain), F(ab)'2 Fragment - Affinity Pure, TRITC Conjugate
Part Number:	GtxHu-001-GRHOX
Concentration:	1.0 mg/ml (E 1% at 280 nm = 13.0)
Amount:	0.5 mg
Conjugate:	Tetramethylrhodamine-5-isothiocyanate (TRITC) Amax = 550 nm; Emax = 570 nm
Form:	Lyophilized
Purification:	Affinity purified using solid phase Human IgA
Purity: Host:	> 90% based on SDS-PAGE Small amounts of intact IgG may be present. Goat
Immunogen:	Purified Human IgA, (α chain)
Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/lgG free
Preservative:	0.05% (w/v) Sodium Azide
Reconstitution:	Rehydrate with 0.55 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh
	working dilution daily.
Storage:	2-8 °C
Shelf Life:	1 year from date of receipt. Prepare working dilution prior to use and then discard.
Specificity:	Based on IEP, this antibody reacts with:
	• heavy (α) chains on human IgA
Cross Reactivity:	Based on IEP, no reactivity is observed to:
	 non-immunoglobulin human serum immunoglobulins light chains on all human immunoglobulins
Country of Origin:	Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.
Applications:	Flow Cytometry
	Immunofluorescence
	ELISA

Immunomicroscopy

Disclaimer:

For *in vitro* Laboratory Use Only. Not for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license under any patent of ImmunoReagents, Inc. **Product may not be resold or modified for resale without prior written approval of ImmunoReagents, Inc.**