## **\* Immuno**Reagents,Inc.

6003 Chapel Hill Road Suite 153 Raleigh, NC 27607

## **Product Specifications**

| Product:<br>Description: | Goat anti-Human IgE (ε chain) - Affinity Pure<br>Goat anti-Human IgE (ε chain) - Affinity Pure, DyLight®650 Conjugate,<br>min x w/bovine, mouse, rabbit serum              |
|--------------------------|--|
| Part Number:             | GtxHu-002-E650NHSX   |
| Concentration:           | 1.0 mg/ml (E 1% at 280 nm = 13.0)  |
| Amount:                  | 1.0 mg   |
| Conjugate:               | DyLight® 650 (Ex = 652 nm; Em = 672 nm)  |
| Form:                    | Lyophilized  |
| Purification:            | Affinity purified using solid phase Human IgE  |
| Purity:                  | Affinity purified antibody is $>$ 95% based on SDS-PAGE  |
| Host:                    | Goat   |
| Immunogen:               | Purified Human IgE, ( $\epsilon$ ) chain   |
| Buffer:                  | 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v)  |
|                          | BSA, Protease/lgG free   |
| Preservative:            | 0.05% (w/v) Sodium Azide   |
| Reconstitution:          | Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room  |
|                          | temperature to dissolve. (Product has been overfilled to ensure complete   |
|                          | recovery.) Centrifuge to remove any particulates. Prepare fresh working  |
| Storego                  | dilution daily.  |
| Storage:<br>Shelf Life:  | Store freeze-dried powder at 2-8 °C.   |
| Shell Lile.              | Product is stable for up to 4 weeks at 2-8 °C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and                               |
|                          | store at -20°C.  |
| Specificity:             | Based on IEP, this antibody reacts with:   |
|                          | ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,   |
|                          | • heavy ( $\epsilon$ ) chains on human IgE   |
| Cross Reactivity:        | Based on IEP, no reactivity is observed to:  |
|                          | <ul> <li>non-immunoglobulin human serum immunoglobulins</li> <li>light chains on all human immunoglobulins</li> <li>serum proteins from bovine, mouse or rabbit</li> </ul> |
| Country of Origin:       | Goat serum was obtained from healthy animals of US origin and under the  |
|                          | care of a registered veterinarian.   |
| Applications:            | Flow Cytometry   |
|                          | Immunofluorescence   |
| Disc. 010 001 0010       |  |

Fax: 919-831-2240

|             | ELISA   |
|-------------|---|
|             | Immunomicroscopy  |
| Disclaimer: | For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not    |
|             | for human or animal consumption. Suggested applications of our products         |
|             | are not recommendations to use our products in violation of any patent or       |
|             | as a license under any patent of ImmunoReagents, Inc. Product may not           |
|             | be resold or modified for resale without prior written approval of              |
|             | ImmunoReagents, Inc.  |
| Trademark:  | DyLight® is a trademark of Thermo Fisher Scientific, Inc. and its subsidiaries. |