*** Immuno**Reagents,Inc.

6003 Chapel Hill Road Suite 153 Raleigh, NC 27607

Product Specifications

Part Number:GtxHu-004-E2RHOXConcentration:1.0 mg/ml (E 1% at 280 nm = 13.0)Amount:1.0 mgConjugate:Tetramethylrhodamine-5-isothiocyanate (TRITC) Amax = 550 nm; Emax = 570 nmForm:LyophilizedPurification:Affinity purified using solid phase Human IgGPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgG, Fc fragmentBuffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Concentration:1.0 mg/ml (E 1% at 280 nm = 13.0)Amount:1.0 mgConjugate:Tetramethylrhodamine-5-isothiocyanate (TRITC) Amax = 550 nm; Emax = 570 nmForm:LyophilizedPurification:Affinity purified using solid phase Human IgGPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgG, Fc fragmentBuffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Amount:1.0 mgConjugate:Tetramethylrhodamine-5-isothiocyanate (TRITC) Amax = 550 nm; Emax = 570 nmForm:LyophilizedPurification:Affinity purified using solid phase Human IgGPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgG, Fc fragmentBuffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Conjugate:Tetramethylrhodamine-5-isothiocyanate (TRITC) Amax = 550 nm; Emax = 570 nmForm:LyophilizedPurification:Affinity purified using solid phase Human IgGPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgG, Fc fragmentBuffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
570 nmForm:LyophilizedPurification:Affinity purified using solid phase Human IgGPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgG, Fc fragmentBuffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Purification:Affinity purified using solid phase Human IgGPurity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgG, Fc fragmentBuffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Purity:Affinity purified antibody is > 95% based on SDS-PAGEHost:GoatImmunogen:Purified Human IgG, Fc fragmentBuffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Host:GoatImmunogen:Purified Human IgG, Fc fragmentBuffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Immunogen:Purified Human IgG, Fc fragmentBuffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Buffer:10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
BSA, Protease/lgG freePreservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Preservative:0.05% (w/v) Sodium AzideReconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
Reconstitution:Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.Storage:2-8 °C
recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Storage: 2-8 °C
dilution daily. Storage: 2-8 °C
Storage: 2-8 °C
5
Shelf Life: 1 year from date of receipt. Prepare working dilution prior to use and then discard.
Specificity: Based on IEP, this antibody reacts with:
 heavy (γ) chains on human IgG
Cross Reactivity: Based on IEP, no reactivity is observed to:
 non-immunoglobulin human serum immunoglobulins light chains on all human immunoglobulins human IgG F(ab) '2 fragment mouse IgG1 serum proteins from bovine, horse, mouse, or rabbit
Country of Origin:
Goat serum was obtained from healthy animals of US origin and under the

Phone: 919-831-2240

Fax: 919-831-2240 email: info@immunoreagents.com

	care of a registered veterinarian.
Applications:	Flow Cytometry
	Immunofluorescence
	ELISA
	Immunomicroscopy
Disclaimer:	For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not
	for human or animal consumption. Suggested applications of our products
	are not recommendations to use our products in violation of any patent or
	as a license under any patent of ImmunoReagents, Inc. Product may not
	be resold or modified for resale without prior written approval of
	ImmunoReagents, Inc.