## **\* Immuno**Reagents,Inc.

6003 Chapel Hill Road Suite 153 Raleigh, NC 27607

## **Product Specifications**

| Product:<br>Description: | Goat anti-Rabbit IgG (H&L) - Affinity Pure<br>Goat anti-Rabbit IgG (H&L) - Affinity Pure, DyLight®405 Conjugate, min x<br>w/bovine, human, mouse IgG/serum |
|--------------------------|--|
| Part Number:             | GtxRb-003-E405NHSX   |
| Concentration:           | 1.0 mg/ml (E 1% at 280 nm = 13.0)  |
| Amount:                  | 1.0 mg   |
| Conjugate:               | DyLight® 405 (Ex = 400 nm; Em = 420 nm)  |
| Form:                    | Lyophilized  |
| Purification:            | Affinity purified using solid phase Rabbit IgG   |
| Purity:                  | Affinity purified antibody is > 95% based on SDS-PAGE  |
| Host:                    | Goat   |
| Immunogen:               | Purified Rabbit IgG, whole molecule  |
| Buffer:                  | 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v)<br>BSA, Protease/lgG free  |
| Preservative:            | 0.05% (w/v) Sodium Azide   |
| Reconstitution:          | Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room  |
|                          | temperature to dissolve. (Product has been overfilled to ensure complete   |
|                          | recovery.) Centrifuge to remove any particulates. Prepare fresh working  |
|                          | dilution daily.  |
| Storage:                 | Store freeze-dried powder at 2-8 °C.   |
| Shelf Life:              | Product is stable for up to 4 weeks at 2-8°C after rehydration. For  |
|                          | extended storage after rehydration, add an equal volume of glycerol and store at -20°C.  |
| Specificity:             | Based on IEP, this antibody reacts with:   |
| Specificity.             | Dased officity, this antibody reacts with.   |
|                          | <ul> <li>heavy (γ) chains on rabbit IgG</li> </ul>   |
|                          | <ul> <li>light chains on all rabbit immunoglobulins</li> </ul>   |
| Cross Reactivity:        | Based on IEP, no reactivity is observed to:  |
|                          | <ul> <li>non-immunoglobulin rabbit serum proteins</li> </ul>   |
|                          | <ul> <li>serum proteins from bovine, human, or mouse</li> </ul>  |
|                          | <ul> <li>IgG from human or mouse</li> </ul>  |
|                          | C C C C C C C C C C C C C C C C C C C  |
| Country of Origin:       | Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.   |
| Applications:            |  |
| Dhama: 010 001 0040      | Flow Cytometry   |
| Phone: 919-831-2240      | Fax: 919-831-2240 email: info@immunoreagents.com   |

|             | Immunofluorescence   |
|-------------|--|
|             | ELISA  |
|             | Immunomicroscopy   |
| Disclaimer: | For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not   |
|             | for human or animal consumption. Suggested applications of our products        |
|             | are not recommendations to use our products in violation of any patent or      |
|             | as a license under any patent of ImmunoReagents, Inc. Product may not          |
|             | be resold or modified for resale without prior written approval of             |
|             | ImmunoReagents, Inc.   |
| Trademark:  | $DyLight^{	extsf{B}}$ is a trademark of Thermo Fisher Scientific, Inc. and its |
|             | subsidiaries.  |