## **\* Immuno**Reagents,Inc.

6003 Chapel Hill Road Suite 153 Raleigh, NC 27607

## **Product Specifications**

Product: Description: Part Number: Concentration: Amount: Conjugate:	Goat anti-Rabbit IgG Fc - Affinity Pure Goat anti-Rabbit IgG Fc - Affinity Pure, TRITC Conjugate GtxRb-004-DRHOX 1.0 mg/ml (E 1% at 280 nm = 13.0) 1.0 mg Tetramethylrhodamine-5-isothiocyanate (TRITC) Amax = 550 nm; Emax = 570 nm
Form:	Lyophilized
Purification:	Affinity purified using solid phase Rabbit IgG
Purity:	Affinity purified antibody is $>$ 95% based on SDS-PAGE
Host:	Goat
Immunogen:	Purified Rabbit IgG Fc
Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/lgG free
Preservative:	0.05% (w/v) Sodium Azide
Reconstitution:	Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.
Storage:	2-8 °C
Shelf Life:	1 year from date of receipt. Prepare working dilution prior to use and then discard.
Specificity:	Based on IEP, this antibody reacts with:
	<ul> <li>heavy (γ) chains on rabbit IgG</li> </ul>
Cross Reactivity:	Based on IEP, no reactivity is observed to:
	<ul> <li>non-immunoglobulin rabbit serum proteins</li> <li>light chains on all rabbit immunoglobulins</li> <li>Rabbit IgG F(ab)'2 fragment</li> </ul>
Country of Origin:	Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.
Applications:	Flow Cytometry
	Immunofluorescence
	ELISA

Immunomicroscopy

Disclaimer:

For *in vitro* Laboratory Use Only. Not for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license under any patent of ImmunoReagents, Inc. **Product may not be resold or modified for resale without prior written approval of ImmunoReagents, Inc.**