## **\* Immuno**Reagents,Inc.

6003 Chapel Hill Road Suite 153 Raleigh, NC 27607

## **Product Specifications**

Product: Description:	Goat anti-Rat IgG (H&L) - Affinity Pure Goat anti-Rat IgG (H&L) - Affinity Pure,FITC Conjugate, min x w/ Bovine, Goat, Hamster, Horse, Human, Mouse, Rabbit IgG/Serum
Part Number: Concentration: Amount:	GtxRt-003-F2FITC 1.50 mg/ml (E 1% at 280 nm = 13.0) 2.0 mg
Conjugate: Form: Purification:	Fluorescein-5-isothiocyanate (FITC) Amax = 494 nm; Emax = 518 nm Clear, fluorescent yellow liquid Affinity purified using solid phase Rat IgG
Purity:	Affinity purified antibody is > 95% based on SDS-PAGE
Host: Immunogen:	Goat Purified Rat IgG, whole molecule
Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free
Preservative:	0.05% (w/v) Sodium Azide
Storage:	2-8 °C
Shelf Life:	1 year from date of receipt. Prepare working dilution prior to use and then discard.
Specificity:	Based on IEP, this antibody reacts with:
	<ul> <li>heavy (γ) chains on rat IgG</li> </ul>
	<ul> <li>light chains on all rat immunoglobulins</li> </ul>
Cross Reactivity:	Based on IEP, no reactivity is observed to:
	<ul> <li>non-immunoglobulin rat serum proteins</li> </ul>
	<ul> <li>IgG from Goat, Hamster, Human, Mouse, or Rabbit</li> </ul>
	<ul> <li>Serum from Bovine, Goat, Horse, Human, Mouse, or Rabbit</li> </ul>
Country of Origin:	Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.
Applications:	Flow Cytometry
	Immunofluorescence
	ELISA
Disoloimor	Immunomicroscopy
Disclaimer:	For <i>in vitro</i> Laboratory Use Only. Not for diagnostic or therapeutic use. Not for human or animal consumption. Suggested applications of our products
	is numar of annual consumption. Suggested applications of our products

are not recommendations to use our products in violation of any patent or as a license under any patent of ImmunoReagents, Inc. **Product may not be resold or modified for resale without prior written approval of ImmunoReagents, Inc.**