

6003 Chapel Hill Road · Suite # 153 · Raleigh, NC 27607

Certificate of Analysis


| Product: | Goat anti-Human IgG Fc, HRP Conjugate |
|--------------------|---|
| Part Number: | GtxHu-004-DHRPX |
| Lot Number: | 90-107-080921 |
| Amount: | 1.0 mg of lyophilized antibody (E $^{1\%}$ at 280 nm = 13.0) |
| | (Product has been overfilled to ensure maximum recovery) |
| Label: | Horseradish Peroxidase (HRP) |
| Form: | Lyophilized |
| Purification: | Affinity purified using solid phase Human IgG |
| Purity: | Affinity purified antibody is \geq 95 % based on SDS-PAGE |
| Host: | Goat |
| Immunogen: | Purified Human IgG Fc |
| Buffer: | 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) |
| | BSA, Protease/IgG free |
| Preservative: | 0.1 % (v/v) ProClin 150 – Use of Sodium Azide will inhibit enzyme activity of |
| | horseradish peroxidase |
| Reconstitution: | Rehydrate with 1.1 ml of deionized water. Allow reconstituted product to stand |
| | at room temperature for at least 30 minutes prior to dilution. If necessary, |
| | centrifuge to remove any particulates. Prepare fresh working dilution daily. |
| Storage: | Store lyophilized material at 2-8 °C. For storage at -20 °C after reconstitution, |
| | dilute with an equal volume of glycerol to prevent loss of enzymatic activity. |
| | Prepare working dilution prior to use and then discard. |
| Shelf Life: | 1 year from date of receipt |
| Specificity: | Based on IEP, this antibody reacts with: |
| | heavy (γ) chains on Human IgG |
| Cross Reactivity: | Based on IEP, no reactivity is observed to: |
| | light chains on all Human immunoglobulins |
| | non-immunoglobulin Human serum proteins |
| Country of Origin: | Goat antiserum was obtained from healthy animals of US origin and under the |
| | care of a registered veterinarian. |
| Note: | This conjugate was prepared using Part # GtxHu-004-D, Lot # 81-86-060821. |
| Applications: | This conjugate is suitable for all immunoassay applications. The optimal |
| | working dilution should be determined by the investigator. Suggested starting |
| | dilution(s): |
| | • 1:500-1:5,000 for Immunohistochemistry |
| | • 1:200-1:5,000 for ELISA/Western blot |
| Disclaimer: | For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not for |
| | human or animal consumption. The proper selection and use of our products is |
| | the sole responsibility of the end user, and therefore we can offer no guarantee to |
| | a specific experimental outcome. Suggested applications of our products are not |
| | recommendations to use our products in violation of any patent or as a license |
| | under any patent of ImmunoReagents, Inc. |
| | Product may not be resold or modified for resale without prior written |
| | approval of ImmunoReagents, Inc. |

*** Immuno**Reagents,Inc.

6003 Chapel Hill Road · Suite # 153 · Raleigh, NC 27607

Testing Results

| PRODUCT: PART NUMBER: LOT NUMBER: | Goat anti-Human IgG Fc, HRP Conjugate GtxHu-004-DHRPX 90-107-080921 | |
|---|--|----------------|
| <u>TESTS</u> | SPECIFICATIONS | <u>RESULTS</u> |
| Appearance: | Reddish powder | PASS |
| Purity: | Two predominant bands at 50 and 24 kDa based on SDS-PAGE | PASS |
| Specificity: | Based on IEP, this antibody has precipitin bands with: heavy (γ) chains on Human IgG | PASS |
| Cross Reactivity: | Based on IEP, no precipitin bands are observed to: light chains on all Human immunoglobulins non-immunoglobulin Human serum proteins | PASS PASS |
| ELISA: | Dilution giving A^{450} reading of ≥ 1.0 when tested against Human IgG (H&L) coated at 2 µg/ml | 1: 331,000 |
| SDS-PAGE | Immunoelectrophoresis (IEP) | |



Date: August 30, 2021

Signature: ____

Quality Control