

6003 Chapel Hill Road · Suite # 153 · Raleigh, NC 27607

Certificate of Analysis

| Product: | Chicken anti-Goat IgG (H&L), HRP Conjugate |
|---|---|
| Part Number: | CkxGt-003-DHRPX |
| Lot Number: | 75-71-030520 |
| Amount: | 1.0 mg of lyophilized antibody (E $^{1\%}$ at 280 nm = 13.2) |
| | (Product has been overfilled to ensure maximum recovery) |
| Label: | Horseradish Peroxidase (HRP) |
| Form: | Lyophilized |
| Purification: | Affinity purified using solid phase Goat IgG |
| Purity: | Affinity purified antibody is \geq 95 % based on SDS-PAGE. |
| Host: | Chicken Dwiffed Cost IsC whole melecule |
| Immunogen: Buffer: | Purified Goat IgG, whole molecule 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, |
| Dullel. | 1 % (w/v) BSA, Protease/IgG free |
| Preservative: | 0.1 % (v/v) ProClin 150– Use of Sodium Azide will inhibit enzyme activity of |
| 110001/001/01 | horseradish peroxidase |
| Reconstitution: | Rehydrate with 1.1 ml of deionized water. Allow reconstituted product to stand at |
| | least 30 minutes at room temperature prior to dilution. If necessary, centrifuge to |
| | remove any particulates. Prepare fresh working dilution daily. |
| Storage: | Store lyophilized material at 2-8 °C. For long term storage after reconstitution, dilute |
| | with 50 % glycerol and store at -20 °C as a liquid. Prepare working dilution prior to |
| 01 101 10 | use and then discard. |
| Shelf Life: | Lyophilized product is stable 1 year from date of receipt. |
| Specificity: | Based on IEP, this antibody reacts with: |
| opeomeny. | |
| specificity. | • heavy (γ) chains on Goat IgG |
| Specificity. | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins |
| Cross Reactivity: | heavy (γ) chains on Goat IgG |
| | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins |
| | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the |
| Cross Reactivity: Country of Origin: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. |
| Cross Reactivity: Country of Origin: Note: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. |
| Cross Reactivity: Country of Origin: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. |
| Cross Reactivity: Country of Origin: Note: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence |
| Cross Reactivity: Country of Origin: Note: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: |
| Cross Reactivity: Country of Origin: Note: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: 1:500-1:5,000 for Immunohistochemistry |
| Cross Reactivity: Country of Origin: Note: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: |
| Cross Reactivity: Country of Origin: Note: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: 1:500-1:5,000 for Immunohistochemistry |
| Cross Reactivity: Country of Origin: Note: Applications: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: 1:500-1:5,000 for Immunohistochemistry 1:200-1:5,000 for ELISA/Western blot For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not for Human or animal consumption. The proper selection and use of our products is |
| Cross Reactivity: Country of Origin: Note: Applications: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: 1:500-1:5,000 for Immunohistochemistry 1:200-1:5,000 for ELISA/Western blot For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not for Human or animal consumption. The proper selection and use of our products is the sole responsibility of the end user, and therefore we can offer no guarantee to |
| Cross Reactivity: Country of Origin: Note: Applications: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: 1:500-1:5,000 for Immunohistochemistry 1:200-1:5,000 for ELISA/Western blot For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not for Human or animal consumption. The proper selection and use of our products is the sole responsibility of the end user, and therefore we can offer no guarantee to a specific experimental outcome. Suggested applications of our products are not |
| Cross Reactivity: Country of Origin: Note: Applications: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: 1:500-1:5,000 for Immunohistochemistry 1:200-1:5,000 for ELISA/Western blot For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not for Human or animal consumption. The proper selection and use of our products is the sole responsibility of the end user, and therefore we can offer no guarantee to a specific experimental outcome. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license |
| Cross Reactivity: Country of Origin: Note: Applications: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: 1:500-1:5,000 for Immunohistochemistry 1:200-1:5,000 for ELISA/Western blot For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not for Human or animal consumption. The proper selection and use of our products is the sole responsibility of the end user, and therefore we can offer no guarantee to a specific experimental outcome. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license under any patent of ImmunoReagents, Inc. |
| Cross Reactivity: Country of Origin: Note: Applications: | heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins Based on IEP, no reactivity is observed to: non-immunoglobulin Goat serum proteins Chicken Ig Fraction was prepared using eggs from hens of US origin and under the care of a registered veterinarian. This product was prepared from Part # CkxGt-003-D, lot # 47-146-020117. The optimal working dilution should be determined by the investigator. Suitable for use in: ELISA, WB, IHC, Chemiluminescence Suggested starting dilutions: 1:500-1:5,000 for Immunohistochemistry 1:200-1:5,000 for ELISA/Western blot For in vitro Laboratory Use Only. Not for diagnostic or therapeutic use. Not for Human or animal consumption. The proper selection and use of our products is the sole responsibility of the end user, and therefore we can offer no guarantee to a specific experimental outcome. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license |

*ImmunoReagents,Inc.

6003 Chapel Hill Road · Suite # 153 · Raleigh, NC 27607

Testing Results

| PRODUCT: PART NUMBER: LOT NUMBER: | Chicken anti-Goat IgG (H&L), HRP Conjugate CkxGt-003-DHRPX 75-71-030520 | | |
|---|--|--------------|--|
| TESTS SPECIFICATIONS RESULTS | | | |
| Appearance: | Reddish powder | PASS | |
| Purity: | Two predominant bands at 60 and 24 kDa based on SDS-PAGE | PASS | |
| Specificity: | Based on IEP, this antibody has precipitin bands with: heavy (γ) chains on Goat IgG light chains on all Goat immunoglobulins | PASS PASS | |
| Cross Reactivity: | Based on IEP, this antibody does not have precipitin bands to:non-immunoglobulin Goat serum proteins | PASS | |
| ELISA: | Dilution giving A^{450} reading of ≥ 1.0 when tested against Goat IgG (H&L) coated at 2 µg/ml | 1: 30,000 | |
| SDS-PAGE | Immunoelectrophoresis (IEP) | | |
| 311 | kDa kDa kDa kDa | 17 | |
| CkxGt-003-D, Lot # 47-146-020117 Mol Weight Standard Date: November 5, 2020 Signature:Quality Control | | | |